

University of Groningen

## Transcriptomic profile of aguR deletion mutant of *Lactococcus lactis* subsp. *cremoris* CECT 8666

Del Rio, Beatriz; Linares, Daniel M; Redruello, Begoña; Martin, Maria Cruz; Fernandez, Maria; de Jong, Anne; Kuipers, Oscar P; Ladero, Victor; Alvarez, Miguel A

*Published in:*  
Genomics Data

*DOI:*  
[10.1016/j.gdata.2015.10.002](https://doi.org/10.1016/j.gdata.2015.10.002)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Del Rio, B., Linares, D. M., Redruello, B., Martin, M. C., Fernandez, M., de Jong, A., Kuipers, O. P., Ladero, V., & Alvarez, M. A. (2015). Transcriptomic profile of aguR deletion mutant of *Lactococcus lactis* subsp. *cremoris* CECT 8666. *Genomics Data*, 6, 228-230. <https://doi.org/10.1016/j.gdata.2015.10.002>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



## Data in Brief

# Transcriptomic profile of *aguR* deletion mutant of *Lactococcus lactis* subsp. *cremoris* CECT 8666

Beatriz del Rio<sup>a</sup>, Daniel M. Linares<sup>a</sup>, Begoña Redruello<sup>a</sup>, Maria Cruz Martin<sup>a</sup>, Maria Fernandez<sup>a</sup>, Anne de Jong<sup>b</sup>, Oscar P. Kuipers<sup>b</sup>, Victor Ladero<sup>a,\*</sup>, Miguel A. Alvarez<sup>a</sup>

<sup>a</sup> Instituto de Productos Lácteos de Asturias, IPLA-CSIC, Paseo Rio Linares s/n, 33300 Villaviciosa, Spain

<sup>b</sup> Department of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands

## ARTICLE INFO

## Article history:

Received 30 September 2015

Accepted 2 October 2015

Available online 9 October 2015

## Keywords:

*Lactococcus lactis*

Putrescine

Agmatine deiminase cluster

*AguR*

Microarray

## ABSTRACT

*Lactococcus lactis* subsp. *cremoris* CECT 8666 (formerly GE2-14) is a dairy strain that catabolizes agmatine (a decarboxylated derivative of arginine) into the biogenic amine putrescine by the agmatine deiminase (AGDI) pathway [1]. The AGDI cluster of *L. lactis* is composed by five genes *aguR*, *aguB*, *aguD*, *aguA* and *aguC*. The last four genes are responsible for the deamination of agmatine to putrescine and are co-transcribed as a single polycistronic mRNA forming the catabolic operon *aguBDAC* [1]. *aguR* encodes a transmembrane protein that functions as a one-component signal transduction system that senses the agmatine concentration of the medium and accordingly regulates the transcription of *aguBDAC* [2], which is also transcriptionally regulated by carbon catabolic repression (CCR) via glucose, but not by other sugars such as lactose and galactose [1,3]. Here we report the transcriptional profiling of the *aguR* gene deletion mutant (*L. lactis* subsp. *cremoris* CECT 8666  $\Delta$ *aguR*) [2] compared to the wild type strain, both grown in M17 medium with galactose as carbon source and supplemented with agmatine. The transcriptional profiling data of *AguR*-regulated genes were deposited in the Gene Expression Omnibus (GEO) database under accession no. GSE59514.

© 2015 Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications	
Organism/cell line/tissue	<i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 (formerly GE2-14)
Sex	N/A
Sequencer or array type	Oligo-based DNA microarray
Data format	Raw and normalized
Experimental factors	<i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 $\Delta$ <i>aguR</i> (test) versus <i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 (reference)
Experimental features	Microarray comparison was performed to identify genes differentially expressed in <i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 $\Delta$ <i>aguR</i> compared to <i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 grown in M17 medium supplemented with 1% galactose (w/v) and 20 mM agmatine.
Consent	N/A
Sample source location	Villaviciosa, Spain

## 1. Direct link to deposited data

Microarray data are accessible in the following link: <http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE59514>.

\* Corresponding author.

## 2. Experimental design, materials and methods

### 2.1. Design of *L. lactis* subsp. *cremoris* CECT 8666 DNA microarrays

*L. lactis* subsp. *cremoris* CECT 8666 DNA microarrays (Agilent Technologies, Santa Clara, CA) were designed using the Agilent eArray (v5.0) program according to the manufacturers' recommendations as described in Linares et al. (2015) [2]. Each microarray (8 × 15 K) was designed to contain spots of two different 60-mer oligonucleotide probes (in duplicate) specific for each of the 2635 coding DNA sequences (CDSs) representing the protein-coding genes of the *L. lactis* subsp. *cremoris* CECT 8666 genome (GenBank accession no. AZSI00000000.1) [4].

### 2.2. Bacterial strains and growth conditions

Table 1 shows the strains used in this study. *L. lactis* subsp. *cremoris* CECT 8666 was originally isolated from a traditional cheese [5]. The mutant strain *L. lactis* subsp. *cremoris* CECT 8666  $\Delta$ *aguR* was constructed by homologous recombination [2]. Both strains were grown in replicates (10 ml each) in M17 medium (Oxoid, Basingstoke, United Kingdom) supplemented with 1% galactose (w/v) and 20 mM agmatine (Sigma-Aldrich, Barcelona, Spain) for 6 h at 30 °C. Cells were harvested by

**Table 1**  
Strains used in this study.

Strain	Description	Reference/source
<i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 <sup>a</sup>	Putrescine producer	[3], CECT <sup>b</sup>
<i>L. lactis</i> subsp. <i>cremoris</i> CECT 8666 $\Delta$ <i>aguR</i>	Non-putrescine producer	[5]

<sup>a</sup> Formerly *L. lactis* subsp. *cremoris* GE2-14.

<sup>b</sup> CECT: Colección Española de Cultivos Tipo.

centrifugation at 8000  $\times$  g for 5 min at 4 °C. The supernatants were removed and cell pellets were frozen in liquid nitrogen and stored at –80 °C.

### 2.3. RNA extraction

RNA extraction was performed as previously described [6] with minor modifications. Briefly, cell pellets were thaw on ice and resuspended in 500  $\mu$ l of TE buffer (10 mM Tris–HCl, 1 mM EDTA pH 8.0) and transferred to screw-capped tubes containing 50  $\mu$ l of 10% SDS, 500  $\mu$ l of phenol:chloroform:isoamyl alcohol (25:24:1) (Sigma-Aldrich), 500 mg of glass beads (75–150  $\mu$ m) (Sigma-Aldrich), and 175  $\mu$ l of Macaloid suspension (Bentone MA, Rheox Inc., Scotland, United Kingdom). Cells were mechanically disrupted in a bead beater at 4 °C. The samples were shaken two times for 45 s. During the shaking intervals the cells were kept on ice for 1 min. The samples were then centrifuged at 8000  $\times$  g for 10 min at 4 °C. The upper phase was transferred to fresh tubes containing 500  $\mu$ l chloroform:isoamyl alcohol (24:1) and centrifuged for 5 min at 4 °C. 500  $\mu$ l of the upper phase was transferred to fresh tubes containing 1 ml of lysis/binding buffer of the High Pure RNA Isolation Kit (Roche Diagnostics GmbH, Mannheim, Germany). All subsequent steps including the DNaseI treatment were performed following the instructions provided by the manufacturer. The concentration and quality of the RNA were checked on a NanoDrop spectrophotometer (Thermo Scientific, Landsmeer, The Netherlands).

### 2.4. Synthesis of cDNA

The synthesis of cDNA was performed using 20  $\mu$ g of total RNA and the SuperScript® III Reverse Transcriptase kit (Life Technologies, Bleiswijk, Netherlands), as described in Shafeeq et al. (2015) [6]. After the cDNA was synthesized, the mRNA of the reverse transcription mixture was denaturalized by adding 3  $\mu$ l of 2.5 mM NaOH for 15 min at 37 °C. The NaOH was neutralized by adding 15  $\mu$ l of 2 M HEPES free acid. The cDNA was purified using the NucleoSpin Gel and PCR Clean-up kit (Macherey-Nager, Landsmeer, The Netherlands). Briefly, 200  $\mu$ l of NTC buffer was mixed with the unpurified cDNA, added to a column and centrifuged for 1 min at 11,000  $\times$  g. The column was washed first with 600  $\mu$ l of buffer NT3 and then with 500  $\mu$ l 80% ethanol. The residual ethanol was completely removed by centrifugation for 2 min at 11,000  $\times$  g. To elute the cDNA, 60  $\mu$ l of 0.1 M sodium bicarbonate pH 9.0 was added to the column and incubated for 1 min at room temperature. Purified cDNA was collected by centrifugation for 1 min at 11,000  $\times$  g and was immediately labeled.

### 2.5. Labeling of cDNA

DyLight 550 NHS ester and DyLight 650 NHS ester (Thermo Scientific) were used to label the cDNAs. Dyes were dissolved in 200  $\mu$ l of DMSO (dimethyl sulfoxide) (Sigma-Aldrich). 60  $\mu$ l of purified cDNA (in 0.1 M sodium bicarbonate pH 9.0, see above) labeled with 5  $\mu$ l DyLight 550 or DyLight 650 in the dark for 90 min at room temperature. Labeled cDNA was purified using NucleoSpin Gel and PCR Clean-up columns as described in the previous section, with the exception that cDNA was eluted with 50  $\mu$ l of elution buffer NE of the NucleoSpin Gel and PCR Clean-up kit.

### 2.6. Hybridization and washing

Nine hundred nanograms of DyLight 550- and DyLight 650-labeled cDNA was mixed and hybridized for 17 h at 60 °C in the *L. lactis* subsp. *cremoris* CECT 8666 DNA microarray using the In situ Hybridization Kit Plus, the Hybridization Gasket Slide and the Agilent G2534A Microarray Hybridization Chamber (Agilent Technologies). After hybridization, slides were washed using appropriate washing buffers as recommended by the manufacturer.

### 2.7. Microarray data analysis

Slides were scanned using a GenePix 4200A Microarray Scanner (Molecular Devices, Sunnyvale, CA) and the images analyzed using GenePix Pro v.6.0 software. Background subtraction and LOWESS (locally weighted scatterplot smoothing) normalization were performed using the standard routines provided by GENOME2D software available at <http://server.molgenrug.nl/index.php/analysis-pipeline>. DNA microarray data were obtained from three independent biological replicates and two technical replicates (including a dye swap). Expression ratios were calculated from the comparison of four spots per gene per microarray (a total of 20 measurements per gene). A gene was considered differentially expressed when a *p* value of at least <0.05 was obtained and the expression fold-change was at least >|0.5|. The microarray data were deposited in Gene Expression Omnibus (GEO) database under the accession no. GSE59514.

## 3. Discussion

In this study, we determined the effect of *aguR* deletion on the transcriptomic profile of *L. lactis* subsp. *cremoris* CECT 8666 grown in M17 supplemented with 1% galactose and 20 mM agmatine. The genes *aguB*, *aguD*, *aguA* and *aguC* coding for the proteins needed for the biosynthesis of putrescine through the AGDI pathway are highly downregulated in the  $\Delta$ *aguR* mutant strain, indicating the role of *AguR* as transcriptional activator of the catalytic AGDI genes that results essential for putrescine biosynthesis [2]. The microarray analysis also reveals the low expression of *aguR* in the CECT 8666 wild-type strain. In addition, other 49 genes were downregulated and 41 upregulated in the  $\Delta$ *aguR* mutant strain compared to the CECT 8666 wild-type strain. Further investigations are required to elucidate the role of *AguR* in the regulation of these genes.

## Acknowledgments

This work was performed with the financial support of the Spanish Ministry of Economy and Competitiveness (AGL2013-45431-R); the Plan for Science, Technology and Innovation 2013–2017 funded by the European Regional Development Fund and the Principality of Asturias (GRUPIN14-137); and the Spanish National Research Council (I-LINK 0380).

## References

- [1] D.M. Linares, B. del Rio, V. Ladero, B. Redruello, M.C. Martin, M. Fernandez, M.A. Alvarez, The putrescine biosynthesis pathway in *Lactococcus lactis* is transcriptionally regulated by carbon catabolic repression, mediated by CcpA. *Int. J. Food Microbiol.* 165 (1) (2013) 43–50, <http://dx.doi.org/10.1016/j.ijfoodmicro.2013.04.021> (43–50).
- [2] D.M. Linares, B. del Rio, B. Redruello, V. Ladero, M.C. Martin, A. de Jong, O.P. Kuipers, M. Fernandez, M.A. Alvarez, *AguR* is a transmembrane transcription activator of the putrescine biosynthesis operon in *Lactococcus lactis*, and acts in response to agmatine concentration. *Appl. Environ. Microbiol.* 81 (18) (2015) 6145–6157, <http://dx.doi.org/10.1128/AEM.00959-15>.
- [3] B. del Rio, V. Ladero, B. Redruello, D.M. Linares, M. Fernandez, M.C. Martin, M.A. Alvarez, Lactose-mediated carbon catabolite repression of putrescine production in dairy *Lactococcus lactis* is strain dependent. *Food Microbiol.* 48 (2015) 163–170, <http://dx.doi.org/10.1016/j.fm.2014.11.018>.
- [4] V. Ladero, B. del Rio, D.M. Linares, M. Fernandez, B. Mayo, M.C. Martin, M.A. Alvarez, Genome sequence analysis of the biogenic amine-producing strain *Lactococcus lactis*

- subsp. *cremoris* CECT 8666 (Formerly GE2-14). Genome Announce 2 (5) (2014), e01088-14, <http://dx.doi.org/10.1128/genomeA.01088-14>.
- [5] V. Ladero, F.P. Rattray, B. Mayo, M.C. Martin, M. Fernandez, M.A. Alvarez, Sequencing and transcriptional analysis of the biosynthesis gene cluster of putrescine-producing *Lactococcus lactis*. Appl. Environ. Microbiol. 77 (18) (2011) 6409–6418, <http://dx.doi.org/10.1128/AEM.05507-11>.
- [6] S. Shafeeq, M. Afzal, B. Henriques-Normark, O.P. Kuipers, Transcriptional profiling of UlaR-regulated genes in *Streptococcus pneumoniae*. Genomics Data 4 (2015) 57–59, <http://dx.doi.org/10.1016/j.gdata.2015.02.004>.